

New fossil diatoms from Ontario. II. *Achnanthes duthii* sp. nov., *Achnanthes undulatus* sp. nov., and *Neidium distincte-punctatum* Hustedt var. *major* var. nov.*

M. R. SREENIVASA

Department of Biology, University of Waterloo, Waterloo, Ontario, Canada.

Accepted December 14, 1970

Three new taxa of diatoms, *Achnanthes duthii* sp. nov., *Achnanthes undulatus* sp. nov., and *Neidium distincte-punctatum* Hustedt var. *major* var. nov., are described from a sediment core obtained from Sunfish Lake, Waterloo County, Ontario.

Phycologia, 10(1): 79-82 (1971)

Introduction

Sunfish Lake, Waterloo County, Ontario is a small meromictic lake of 8.3 ha in area and a maximum depth of 20 m. Its limnology has been described by Duthie and Carter (1970). The surface waters are moderately alkaline with about 125 mg/l CaCO₃ and a pH of about 8.5. Because of the pronounced concavity of the lake basin, the littoral region is very narrow but dense patches of *Chara* spp. occur in places.

A sediment core of nearly 5 m in length was obtained for paleolimnological investigation. Over 300 taxa of diatoms have been identified from the core (Sreenivasa, 1970) and three taxa, described herein, proved to be new to science. *Achnanthes duthii* sp. nov. and *Achnanthes undulatus* sp. nov. occurred in low numbers in samples between 30 cm and 70 cm from the top of the core. *Neidium distincte-punctatum* Hustedt var. *major* var. nov. occurred in large numbers at a depth of 320 cm in the core. None of the new taxa have been found in recent collections. Paleolimnology, diatom sequence of the core, and its interpretation is in preparation.

***Achnanthes duthii* sp. nov.**

Valva elliptico-lanceolata vel linearis-lanceolata; extremitas obtuse-rotundatis. Rapho-valva: area axialis plane lanceolata, raphe tenui, extremitas proximalibus leviter dilatatis et extremitas distalibus unilateraliter flectis. Valva sine raphe: pseudo-raphe anguste linearis-




FIG. 1. A-D, *Achnanthes duthii* sp. nov. E, *Achnanthes undulatus* sp. nov.

* From Ph.D. thesis, University of Waterloo, Waterloo, Ontario, 1970.

lanceolata; area centrali leviter dilatata et extimus distalibus unilateraliter flectis. Longitudo 20–33 μ ; latitudo 4.4–5 μ . Striis 20–24/10 μ , leviter radialibus vel parallelis, leviter latioribus et longius distantibus ad centrum quam alibi (Fig. 1 A–D; Fig. 2 A–C). Our slide number *SFL 5*. HOLOTYPE: accession number *CANA 8408*, National Museum of Canada, Ottawa. LOCALITY: Waterloo; Sunfish Lake. COLLECTORS: H. R. N. Eydt and H. C. Duthie, 1967.

Valve elliptical lanceolate to linear lanceolate, ends obtusely rounded. Raphe valve: axial area of raphe valve distinctly lanceolate, raphe




FIG. 2. A–C, *Achnanthes duthii* sp. nov. D, *Achnanthes undulatus* sp. nov.




FIG. 3. A–C, *Neidium disticto-punctatum* Hustedt var. major var. nov.

thin, proximal ends slightly enlarged, distal ends unilaterally bent. Rapheless valve: pseudo raphe narrowly linear lanceolate slightly widened at the central area and unilaterally bent at the distal ends. Length 20–33 μ , width 4.4–5 μ , striae 20–24/10 μ . Striae slightly radial to parallel; slightly broader and farther apart at the center than elsewhere.

This taxon resembles *Achnanthes nollii* (Bock, *Nachr. Naturw. Mus. Aschaffenburg*, 38: 54–56, pl. 5, fig. 7–9, 1953); it differs in that the axial area is more lanceolate rather than linear and the terminal raphe pores are unilaterally bent rather than straight. This taxon is slightly larger in size and the striae are far fewer in number than in *A. nollii* (see Patrick, R. and Reimer, C. W., 1966, p. 253, pl. 16, fig. 7–8).

This taxon is very similar to *Achnanthes lata* Hust. in general appearance, but differs from it in its slightly larger size; in the number of striae, which are far less; the median striae are thicker and far apart; and in the absence of irregular longitudinal lines crossing the striae.

Achnanthes

Valva linearis-latis; apicibus proparatis a valva. In centrali longitudine proximalibus extensis. Valva sine raphe centrali longitudine leviter lato. Longitudo 23–25/10 μ , plus (Fig. 1 E, 2 D). accession number Canada, Ottawa. COLLECTORS: H.

Valve linearis-late sides. Ends proparate set off from area linear, cuneangular. Raphe slightly bulbous, distal valve: axial area longitudinally narrow. Length 23–25/10 μ , plus in both the valves.

This taxon does not agree with any species of *Achnanthes*.

Achnanthes sp. nov. occurs from the top of zone *C₃* of the Fagus. Dominant are *Cyclotella* and *Stephanodiscus* assemblages at be alkaline and

Neidium

Valvae ellipticas-rostratis, obtuse leviter decrecentes. Extremis proximalibus praebentibus una raphe-axe. Area proximalia et transversa bifurcatis et transverse ellipticas leviter radialibus et convergentibus 9–11, 63–90 μ , latitudine. Our slide number CANA 8408. tawa. LOCALITY: TORS: H. R. N. E.

Valves ellipticas rounded sub-subsquadrangular, slightly oblique.

***Achnanthes undulatus* sp. nov.**

Valva linearis-lanceolata, lateribus leviter triundulatis; apicibus protractis, subrostratis vel rostratis disparatis a valva. Rapho-valva: area axiali linearis, area centrali longitudinaliter rectangularis, raphe tenuis, proximalibus extremis leviter bulbosis, distalibus rectis. Valva sine raphe: area axiali linearis, recta; area centrali longitudinaliter rectangularis, et extremo distali leviter lato. Longitudo 23.5 μ , latitudo 4.4 μ , striae 23–25/10 μ , plus minusve parallelis in utraque valva (Fig. 1 E, 2 D). Our slide number SFL 3. HOLOTYPE: accession number CANA 8409, National Museum of Canada, Ottawa. LOCALITY: Waterloo; Sunfish Lake. COLLECTORS: H. R. N. Edyt and H. C. Duthie, 1967.

Valve linear lanceolate with slightly triundulate sides. Ends protracted, subrostrate to rostrate set off from the valve. Raphe valve: axial area linear, central area longitudinally rectangular. Raphe fine with proximal ends slightly bulbous, distal ends straight. Rapheless valve: axial area linear straight, central area longitudinally rectangular, distal ends slightly broad. Length 23.5 μ , breadth 4.4 μ , striae 23–25/10 μ , more or less parallel throughout in both the valves.

This taxon does not seem to have any similarity with any species described.

Achnanthes duthii sp. nov. and *A. undulatus* sp. nov. occurred between 30 cm and 70 cm from the top of the core, which is related to zone *C₃* of the pollen sequence (*Tsuga* and *Fagus*). Dominant species associated with them are *Cyclotella glomerata*, *C. kutziniana*, and *Stephanodiscus hantzschii*. From the overall assemblages at these levels the water seems to be alkaline and mesotrophic to eutrophic.

***Neidium distincte-punctatum* Hust. var. *major* var. nov.**

Valvae elliptico-lanceolatae, extremis subrostratis vel rostratis, obtuse rotundatis. Raphe leviter obliqua, leviter decrecenti prope extrema distalia et proximalia. Extremis proximalibus diverse cravatis, nonnunquam praebentibus uncis proximales in raphibus, tenuiores rapho-axe. Area axiali leviter angustiore ad extrema proximalia et distalia. Distalibus extremis raphis bifurcatis et tenioribus axe raphis. Area centrali transverse elliptica vel paene quadrangulata. Striae leviter radialibus et ad extremitates parallelis vel convergentibus 9–11/10 μ ; puncta 12–14/10 μ , longitudine 63–90 μ , latitudo 19–24 μ (Fig. 3 A–C, Fig. 4). Our slide number SFL 32. HOLOTYPE: accession number CANA 8410, National Museum of Canada, Ottawa. LOCALITY: Waterloo; Sunfish Lake. COLLECTORS: H. R. N. Edyt and H. C. Duthie, 1967.

Valves elliptical lanceolate with broad, bluntly rounded subrostrate to rostrate ends. Raphe slightly oblique, narrowing slightly near the

distal and proximal ends. Proximal ends curved in opposite direction, some show proximal raphe hooks; these are thinner than the raphe axis. Distal raphe ends diforcate, also thinner than raphe axis. Axial area slightly narrower at the distal and proximal ends, central area transversely elliptical to nearly quadrangular. Striae slightly radial becoming parallel to convergent at the very ends. Length, 63–90 μ , width 14–24 μ , striae 9–11/10 μ , puncta 12–14/10 μ .

This taxon is closely related to *Neidium distincte-punctatum* Hustedt, but is larger and the valve ends are broader. Hustedt gives length, 40–45 μ , width 16 μ , striae 11/12–10 μ (Hustedt, 1930, p. 247, fig. 386).

The illustration of *Neidium distincte-punctatum* Hustedt in Stoermer (1963) reveals a length of 60 μ , width 18 μ , striae 12/10 μ .




FIG. 4. A–F, *Neidium distincte-punctatum* Hustedt var. *major* var. nov.

It is possible that these are the larger forms where the proximal ends of the raphe are curved in opposite direction and intergrade to form raphe hooks. The shape of the valve ends intergrade from the types given by Hustedt and Stoermer to broad bluntly rounded rostrate to subrostrate ends.

These observations indicate that this is a larger variety of *Neidium distincte-punctatum* Hustedt.

Neidium distincte-punctatum Hustedt var. nov., was found in the core at a depth of 320 cm from the surface, which corresponds with the zone B (pine) of the postglacial.

The dominant species occurring in association are *Cyclotella bodanica* and *Fragilana lapponica*. The overall assemblage indicates deposition in indifferent, oligotrophic water.

Acknowledgements

The assistance of Dr. Ruth Patrick and Dr. Charles W. Reimer of the Academy of Natural Sciences of Philadelphia in confirming the

new taxa is gratefully acknowledged. I wish to thank Dr. P. Keresztes of the University of Waterloo for correcting the Latin diagnoses. The work was supported by a grant to Dr. H. C. Duthie, University of Waterloo, from the Department of Energy, Mines and Resources, Ottawa.

References

- DUTHIE, H. C., AND CARTER, J. C. H. 1970. The meromixis of Sunfish Lake, Southern Ontario. *J. Fish. Res. Bd. Can.*, 27: 847-856.
- HUSTEDT, F. 1930. Bacillariophyta (Diatomeae). In Pascher, A. [ed.] *Die Süßwasserflora Mitteleuropas*. Gustav Fisher, Jena.
- PATRICK, R., AND REIMER, C. W. 1966. *The Diatoms of the United States*. Monogr. 13, Acad. Nat. Sci. Philadelphia, 688 p.
- SREENIVASA, M. R. 1970. *Recent and Extant Diatom Assemblages in Southern Ontario*. Ph.D. thesis, University of Waterloo, 450 p.
- STOERMER, E. F. 1963. New taxa and new United States records of the diatom genus *Neidium* from west Lake Okovoji, Iowa. *Notulae Naturae*, 358: 1-9. Acad. Nat. Sci. Philadelphia.